

CORRESPONDENCE

Severity of SARS-CoV-2 Reinfections as Compared with Primary Infections

TO THE EDITOR: Qatar had a first wave of infections with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from March through June 2020, after which approximately 40% of the population had detectable antibodies against SARS-CoV-2. The country subsequently had two back-to-back waves from January through May 2021, triggered by the introduction of the B.1.1.7 (or alpha) and B.1.351 (or beta) variants.¹ This created an epidemiologic opportunity to assess reinfections.

Using national, federated databases that have captured all SARS-CoV-2–related data since the onset of the pandemic (Section S1 in the Supplementary Appendix, available with the full text of this letter at NEJM.org), we investigated the risk of severe disease (leading to acute care hospitalization), critical disease (leading to hospitalization in an intensive care unit [ICU]), and fatal disease caused by reinfections as compared with primary infections in the national cohort of 353,326 persons with polymerase-chain-reaction (PCR)–confirmed infection between February 28, 2020, and April 28, 2021, after exclusion of 87,547 persons with a vaccination record. Primary

infection was defined as the first PCR-positive swab. Reinfection was defined as the first PCR-positive swab obtained at least 90 days after the primary infection. Persons with reinfection were matched to those with primary infection in a 1:5 ratio according to sex, 5-year age group, nationality, and calendar week of the PCR test date (Fig. S1 and Table S1 in the Supplementary Appendix). Classification of severe, critical, and fatal Covid-19 followed World Health Organization guidelines, and assessments were made by trained medical personnel through individual chart reviews.

Of 1304 identified reinfections, 413 (31.7%) were caused by the B.1.351 variant, 57 (4.4%) by the B.1.1.7 variant, 213 (16.3%) by “wild-type” virus, and 621 (47.6%) were of unknown status (Section S1 in the Supplementary Appendix). For reinfected persons, the median time between first infection and reinfection was 277 days (interquartile range, 179 to 315). The odds of severe disease at reinfection were 0.12 times (95% confidence interval [CI], 0.03 to 0.31) that at primary infection (Table 1). There were no cases of critical disease at reinfection and 28 cases at primary infection (Table S3), for an odds ratio of 0.00

Table 1. Severity of SARS-CoV-2 Reinfections as Compared with Primary Infections in the Population of Qatar.

Disease Outcome*	Reinfection†	Primary Infection†	Odds Ratio (95% CI)
	<i>no. of persons with outcome/no. of persons with infection that was not severe, critical, or fatal</i>		
Severe disease	4/1300	158/6095	0.12 (0.03–0.31)
Critical disease	0/1300	28/6095	0.00 (0.00–0.64)
Fatal disease	0/1300	7/6095	0.00 (0.00–2.57)
Severe, critical, or fatal disease	4/1300	193/6095	0.10 (0.03–0.25)

* Severe disease, critical disease, and fatal disease were defined on the basis of the World Health Organization criteria for classifying severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection severity and coronavirus disease 2019 (Covid-19)–related death.

† Reinfections were matched with up to five primary infections according to sex, 5-year age group, nationality, and calendar week of polymerase-chain-reaction testing. The final sample therefore includes persons with reinfection who were matched to five or fewer persons with primary infection.

(95% CI, 0.00 to 0.64). There were no cases of death from Covid-19 at reinfection and 7 cases at primary infection, resulting in an odds ratio of 0.00 (95% CI, 0.00 to 2.57). The odds of the composite outcome of severe, critical, or fatal disease at reinfection were 0.10 times (95% CI, 0.03 to 0.25) that at primary infection. Sensitivity analyses were consistent with these results (Table S2).

Reinfections had 90% lower odds of resulting in hospitalization or death than primary infections. Four reinfections were severe enough to lead to acute care hospitalization. None led to hospitalization in an ICU, and none ended in death. Reinfections were rare and were generally mild, perhaps because of the primed immune system after primary infection.

In earlier studies, we assessed the efficacy of previous natural infection as protection against reinfection with SARS-CoV-2^{2,3} as being 85% or greater. Accordingly, for a person who has already had a primary infection, the risk of having a severe reinfection is only approximately 1% of the risk of a previously uninfected person having a severe primary infection. It needs to be determined whether such protection against severe disease at reinfection lasts for a longer period, analogous to the immunity that develops against other seasonal “common-cold” coronaviruses,⁴ which elicit short-term immunity against mild reinfection but longer-term immunity against more severe illness with reinfection. If this were the case with SARS-CoV-2, the virus (or at least the vari-

ants studied to date) could adopt a more benign pattern of infection when it becomes endemic.⁴

Laith J. Abu-Raddad, Ph.D.

Hiam Chemaitelly, M.Sc.

Weill Cornell Medicine–Qatar

Doha, Qatar

lja2002@qatar-med.cornell.edu

Roberto Bertollini, M.D., M.P.H.

Ministry of Public Health

Doha, Qatar

for the National Study Group for COVID-19
Epidemiology*

*Members of the National Study Group for COVID-19 Epidemiology are listed in the Supplementary Appendix, available with the full text of this letter at NEJM.org.

Supported by the Biomedical Research Program and the Biostatistics, Epidemiology, and Biomathematics Research Core at Weill Cornell Medicine–Qatar; the Ministry of Public Health; Hamad Medical Corporation; and Sidra Medicine. The Qatar Genome Program supported the viral genome sequencing.

Disclosure forms provided by the authors are available with the full text of this letter at NEJM.org.

This letter was published on November 24, 2021, at NEJM.org.

1. Abu-Raddad LJ, Chemaitelly H, Butt AA. Effectiveness of the BNT162b2 Covid-19 vaccine against the B.1.1.7 and B.1.351 variants. *N Engl J Med* 2021;385:187-9.
2. Abu-Raddad LJ, Chemaitelly H, Coyle P, et al. SARS-CoV-2 antibody-positivity protects against reinfection for at least seven months with 95% efficacy. *EClinicalMedicine* 2021;35:100861.
3. Abu-Raddad LJ, Chemaitelly H, Malek JA, et al. Assessment of the risk of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) reinfection in an intense reexposure setting. *Clin Infect Dis* 2021;73(7):e1830-e1840.
4. Lavine JS, Bjornstad ON, Antia R. Immunological characteristics govern the transition of COVID-19 to endemicity. *Science* 2021;371:741-5.

DOI: 10.1056/NEJMc2108120

Correspondence Copyright © 2021 Massachusetts Medical Society.